

THE SEQIRUS PORTFOLIO OF INFLUENZA VACCINES—2021-2022 SEASON



Seqirus offers a wide range of influenza vaccines for patients 6 months to 65 years of age and older.

PACKAGING	AGES	PRESENTATION	DOSING	CPT CODE	CARTON NDC #	SYRINGE/VIAL LABEL NDC #	PRODUCT DESCRIPTION	CVX CODE [§]	MVX CODE
	65 years and older ¹	10 x 0.5-mL pre-filled syringes* ¹	1 dose	90694 (covered by Medicare Part B)	70461-121-03	70461-321-04	Quadrivalent influenza vaccine manufactured using adjuvant technology ¹	205	SEQ
	2 years and older ²	5-mL multi-dose vial ²	For children 2 through 8 years, 1-2 doses depending on vaccination history	90756	70461-421-10	70461-421-11	Quadrivalent influenza vaccine manufactured with cell-based technology ²	186	SEQ
		10 x 0.5-mL pre-filled syringes* ²		90674	70461-321-03	70461-321-04		171	SEQ
	6 months and older ³	5-mL multi-dose vial ³	For children 6 months through 8 years, 1-2 doses depending on vaccination history For persons 9 years of age and older, 1 dose	90687 (0.25-mL dose)	33332-421-10	33332-421-11	Quadrivalent influenza vaccine manufactured with egg-based technology ³	158	SEQ
	36 months and older ³	10 x 0.5-mL pre-filled syringes* ³		90688 (0.5-mL dose)					
	6 months through 35 months ³	10 x 0.25-mL pre-filled syringes* ³		90686	33332-321-01	33332-321-02		150	SEQ
				90685	33332-221-20	33332-221-21	161	SEQ	

CPT=Current Procedural Terminology; NDC=National Drug Code

*The pre-filled syringes contain no preservative.

†The number of needle punctures should not exceed 20 per multi-dose vial.

§The CVX, or vaccine administered code, indicates which product was used and is used in combination with the manufacturer (MVX) code.

Visit flu.seqirus.com for more information and to place your order.

Please see Important Safety Information on next page and full US Prescribing Information for FLUAD® QUADRIVALENT, FLUCELVAX® QUADRIVALENT, and AFLURIA® QUADRIVALENT.

References: 1. FLUAD QUADRIVALENT. Package insert. Seqirus Inc; 2021. 2. FLUCELVAX QUADRIVALENT. Package insert. Seqirus USA Inc; 2021. 3. AFLURIA QUADRIVALENT. Package insert. Seqirus USA Inc; 2021.

For US Healthcare Professional Use Only
This information does not constitute a guarantee or warranty of coverage benefits or reimbursement

Questions?



Call flu360 | Support
(855) 358-8966, option 2



FLUAD® QUADRIVALENT (Influenza Vaccine, Adjuvanted), AFLURIA® QUADRIVALENT (Influenza Vaccine), and FLUCELVAX® QUADRIVALENT (Influenza Vaccine)

IMPORTANT SAFETY INFORMATION

INDICATIONS AND USAGE

FLUAD® QUADRIVALENT (Influenza Vaccine, Adjuvanted) is an inactivated influenza vaccine indicated for active immunization against influenza disease caused by influenza virus subtypes A and types B contained in the vaccine. FLUAD QUADRIVALENT is approved for use in persons 65 years of age and older. This indication is approved under accelerated approval based on the immune response elicited by FLUAD QUADRIVALENT.

AFLURIA® QUADRIVALENT (Influenza Vaccine) is an inactivated influenza vaccine indicated for active immunization against influenza disease caused by influenza A subtype viruses and type B viruses contained in the vaccine. AFLURIA QUADRIVALENT is approved for use in persons 6 months of age and older.

FLUCELVAX® QUADRIVALENT (Influenza Vaccine) is an inactivated vaccine indicated for active immunization for the prevention of influenza disease caused by influenza virus subtypes A and types B contained in the vaccine. FLUCELVAX QUADRIVALENT is approved for use in persons 2 years of age and older.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

Do not administer FLUAD QUADRIVALENT or AFLURIA QUADRIVALENT to anyone with a history of severe allergic reaction (e.g. anaphylaxis) to any component of the vaccine, including egg protein, or to a previous influenza vaccine. Do not administer FLUCELVAX QUADRIVALENT to anyone with a history of severe allergic reactions (e.g. anaphylaxis) to any component of the vaccine.

WARNINGS AND PRECAUTIONS

If Guillain-Barré syndrome (GBS) has occurred within 6 weeks of receipt of prior influenza vaccine, the decision to give FLUAD QUADRIVALENT, AFLURIA QUADRIVALENT or FLUCELVAX QUADRIVALENT should be based on careful consideration of the potential benefits and risks.

Appropriate medical treatment and supervision must be available to manage possible anaphylactic reactions following administration of the vaccine.

Syncope (fainting) may occur in association with administration of injectable vaccines including FLUAD QUADRIVALENT and FLUCELVAX QUADRIVALENT. Syncope can be accompanied by transient neurological signs such as visual disturbance, paresthesia, and tonic-clonic limb movements. Ensure procedures are in place to avoid falling injury and to restore cerebral perfusion following syncope by maintaining a supine or Trendelenburg position.

The immune response to FLUAD QUADRIVALENT, AFLURIA QUADRIVALENT and FLUCELVAX QUADRIVALENT in immunocompromised persons, including individuals receiving immunosuppressive therapy, may be lower than in immunocompetent individuals.

Vaccination with FLUAD QUADRIVALENT, AFLURIA QUADRIVALENT or FLUCELVAX QUADRIVALENT may not protect all vaccine recipients against influenza disease.

ADVERSE REACTIONS

FLUAD QUADRIVALENT:

The most common (≥ 10%) local and systemic reactions with FLUAD QUADRIVALENT in elderly subjects 65 years of age and older were injection site pain (16.3%), headache (10.8%) and fatigue (10.5%).

AFLURIA QUADRIVALENT:

AFLURIA QUADRIVALENT administered by needle and syringe:

In adults 18 through 64 years, the most commonly reported injection-site adverse reaction was pain (≥ 40%). The most common systemic adverse events were myalgia and headache (≥ 20%).

In adults 65 years of age and older, the most commonly reported injection-site adverse reaction was pain (≥ 20%). The most common systemic adverse event was myalgia (≥ 10%). In children 5 through 8 years, the most commonly reported injection-site adverse reactions were pain (≥ 50%), redness and swelling (≥ 10%). The most common systemic adverse event was headache (≥ 10%).

In children 9 through 17 years, the most commonly reported injection-site adverse reactions were pain (≥ 50%), redness and swelling (≥ 10%). The most common systemic adverse events were headache, myalgia, and malaise and fatigue (≥ 10%).

In children 6 months through 35 months of age, the most commonly reported injection-site reactions were pain and redness (≥ 20%). The most common systemic adverse events were irritability (≥ 30%), diarrhea and loss of appetite (≥ 20%).

In children 36 through 59 months of age, the most commonly reported injection site reactions were pain (≥ 30%) and redness (≥ 20%). The most commonly reported systemic adverse events were malaise and fatigue, and diarrhea (≥ 10%).

The safety experience with AFLURIA (trivalent formulation) is relevant to AFLURIA QUADRIVALENT because both vaccines are manufactured using the same process and have overlapping compositions:

In adults 18 through 64 years of age, the most commonly reported injection-site adverse reactions with AFLURIA (trivalent formulation) when administered by the PharmaJet Stratis Needle-Free Injection System were tenderness (≥ 80%), swelling, pain, redness (≥ 60%), itching (≥ 20%) and bruising (≥ 10%). The most common systemic adverse events were myalgia, malaise (≥ 30%), and headache (≥ 20%).

FLUCELVAX QUADRIVALENT:

In adults 18 through 64 years of age who received FLUCELVAX QUADRIVALENT, the most commonly reported injection-site adverse reactions were pain (≥ 40%), erythema and induration (≥ 10%). The most common systemic adverse events were headache, fatigue and myalgia (≥ 10%).

In adults ≥ 65 years of age who received FLUCELVAX QUADRIVALENT, the most commonly reported injection-site adverse reactions were pain (≥ 20%) and erythema (≥ 10%).

In children 2 through 8 years of age who received FLUCELVAX QUADRIVALENT, the most commonly reported injection-site adverse reactions were tenderness (28.7%), pain (27.9%) and erythema (21.3%), induration (14.9%) and ecchymosis (10.0%). The most common systemic adverse events were sleepiness (14.9%), headache (13.8%), fatigue (13.8%), irritability (13.8%) and loss of appetite (10.6%).

In children and adolescents 9 through 17 years of age who received FLUCELVAX QUADRIVALENT, the most commonly reported injection-site adverse reactions were injection site pain (21.7%), erythema (17.2%) and induration (10.5%). The most common systemic adverse events were headache (18.1%) and fatigue (17.0%).

To report SUSPECTED ADVERSE REACTIONS, contact Seqirus at 1-855-358-8966 or VAERS at 1-800-822-7967 or www.vaers.hhs.gov.

Before administration, please see the full US Prescribing Information for [FLUAD QUADRIVALENT](#), [AFLURIA QUADRIVALENT](#) and [FLUCELVAX QUADRIVALENT](#).

FLUAD® QUADRIVALENT, AFLURIA® QUADRIVALENT and FLUCELVAX® QUADRIVALENT are registered trademarks of Seqirus UK Limited or its affiliates.

PharmaJet® and STRATIS® are registered trademarks of PharmaJet.